Patient & Family Guide 2023 Botox[®] Injection

Register using a self check-in kiosk in the main entrance of the Centennial or Dickson buildings, Victoria General site.

Next, go to the Eye Care Centre in the Centennial building, 2A.



Aussi disponible en français : Injection de Botox[®] (FF85-1783)

Botox® Injection

Your doctor has recommended Botox[®] injections for your blepharospasm (rapid eye blinking) or facial muscle spasms.

What is Botox[®] and how does it work?

- Botox[®] is a purified (germ-free) toxin that can stop muscle spasms.
- Your nerves release chemicals that cause the spasms in your muscles. Botox[®] blocks these chemicals and stops the spasms.

How long do the Botox[®] effects last?

- Botox[®] is not a cure. It helps relieve your symptoms for about 3 months.
- You should see improvement in 3 to 5 days. This will be different for each person and depends on the amount of Botox[®] given.
- If your spasms do not get better, you may need another injection.

On the day of your appointment:

• **Do not** wear any makeup on your eyes or face.

- Go to the main entrance of the Centennial or Dickson building, Victoria General site. Register using a self check-in kiosk.
- Go to the Eye Care Centre in the Centennial building, 2A.
- Plan to be at the Centre for 1 to 1 ½ hours.

During treatment

- You will be asked to sit in a chair with your head well-supported.
- The eye doctor will use a needle to inject the Botox[®] around your eyes and face.

After treatment

- The amount of Botox[®] you will have during this treatment is very small. It does **not** spread to any other parts in your body. You can go back to your usual activities right away.
- **Do not** rub the injection areas after the treatment.

Are there any side effects?

- You may have some of these side effects:
 - › Your upper eyelid may droop.

- > Your eyelid may be puffy.
- Your eye may be dry. You may need eye drops. You **do not** need a prescription for the eye drops.
- You may have trouble closing your upper eyelid.
- > Your eye may be sensitive to light.
- Your eye may tear up because your lower eyelid is weak. This will go away over time as the Botox[®] wears off.
- Your lips may move differently. This may make it hard to talk.
- You may have bruising around the injection areas.
- Double vision (seeing 2 images when you are looking at one object).

These side effects are temporary (will go away over time).

Nova Scotia Health promotes a smoke-free, vape-free, and scent-free environment. Please do not use perfumed products. Thank you! www.nshealth.ca

> Prepared by: Eye Care Centre Designed by: Nova Scotia Health Library Services

The information in this pamphlet is for informational and educational purposes only. The information is not intended to be and does not constitute health care or medical advice. If you have any questions, please ask your health care provider.

WW85-0282 © June 2023 Nova Scotia Health Authority The information in this pamphlet is to be updated every 3 years or as needed.

