

Burn Care


Positioning for Lower Extremity Swelling

Instructions

- ✓ After a burn injury, swelling in the limbs is common. This swelling can be painful and affect your ability to do daily activities.
- ✓ One of the easiest ways to manage swelling is by raising the swollen limb.
- ✓ The goal is to raise the swollen limb slightly above the level of your heart. This helps the extra fluid move back towards your heart to circulate it to the rest of your body.

Positioning in bed

- Lying in bed with your legs raised is the best position to help lower swelling.
- It is best to lie on your back.
- Raise your legs above the level of your heart, while keeping your upper body flat.
- If you have a hospital bed, you can do this by raising the foot of the bed.
- If you are in a regular bed, use pillows to raise each leg. Place the pillows lengthwise under each leg. Use as many pillows as needed to lift your legs above the level of your heart.



Positioning when sitting

- When sitting in a chair, use a stool to raise your legs. You can place pillows on top of the stool if needed.



Positioning when standing

- Walking activates your muscles and helps get fluid moving.
- Avoid standing still for a long time, as this will increase swelling. Try walking around or marching on the spot. While you are standing, move your toes up and down to keep the fluid moving.
- You may use a compression bandage, stockings, or a pressure garment when standing or walking to lower fluid build-up in your ankles. Talk to your occupational therapist or physiotherapist about which type of compression garment or bandage would be best for you.
- After walking or standing for a period of time, return to a chair or bed and raise your legs as described above.

What are your questions? Please ask.

Therapist: _____

Phone: _____

*Prepared by: Rehabilitation Services
Designed by: NSHA Library Services*