

Dobutamine[®] Stress Echocardiogram

Dobutamine® Stress Echocardiogram

What is a Dobutamine® stress echocardiogram (DSE)?

- Your coronary arteries supply blood to your heart. A DSE is a test to find out if there are any blockages in these arteries and, if so, how severe (very bad) the blockages are.
- This test may be done when the usual tests for this problem cannot be done, or have not given clear results.

Who will do the test?

- The test will be done by a cardiologist (heart doctor) and a cardiac sonographer (heart ultrasound specialist). The cardiologist will talk about the test with you, including possible side effects.

How do I get ready for the test?

- There is no preparation for this test.
- Please bring a list of all of your current medications.

- You may want to arrange for a drive home before your test. You can drive yourself home, but you may feel tired after the test.

If you are an outpatient:

- Go to the Halifax Infirmary, Summer Street entrance.
 - › Register at the kiosk on Level 4.
 - › Then, go to the Minor Procedures Clinic on Level 4 to get ready for your test.

What will happen during the test?

- You will be asked to lie on your side.
- Small stickers (electrodes) will be placed on your chest, arms, and legs. The electrodes will then be attached to an echocardiogram machine and an ultrasound machine.
- An intravenous (I.V.) will be inserted (put in) in your hand or arm. The medication Dobutamine® will be given through the I.V. This medication has been given to people with heart problems for many years.
- Your blood pressure and heart rate will be checked every 3 minutes.
- There will always be a doctor present during the test.

- A small device called an ultrasound probe will be moved over your chest. This will give images of your heart.
- Dobutamine® can cause:
 - › Chest pain
 - › Feeling that your heart is pounding fast
 - › Headache
 - › Nausea (upset stomach)
 - › Anxiety
 - › Shakiness
- Your blood pressure may drop a bit or you may have an irregular heart rhythm.
These problems will stop as soon as the Dobutamine® is stopped.
- There are no long-term effects from this medication.
- The cardiologist and cardiac sonographer will monitor the effect of the Dobutamine® on your heart using the electrocardiogram machine.
- You may be given a medication (called a contrast) by I.V. to help make the ultrasound more clear.
- You may be given a medication called atropine by I.V. to raise your heart rate during the test. This medication may make your mouth feel dry.

Will I have any discomfort?

- The test is not painful.
- The I.V. needle may cause temporary discomfort when it is inserted. There may be a bruise where the needle enters the skin.

How long will the test take?

The test will take about 1½ to 2 hours.

After the test

- You will be taken back to the Minor Procedures Clinic recovery area. You will be monitored there for about 1 hour.

You may still feel tired after the recovery time. You are allowed to drive yourself home, but you may want to arrange for someone to pick you up.

[illegible]

This pamphlet is for educational purposes only. It is not intended to replace the advice or professional judgment of a health care provider. The information may not apply to all situations. If you have any questions, please ask your health care provider.

Find all patient education resources here:
www.nshealth.ca/patient-education-resources

Connect with a registered nurse in Nova Scotia any time:
Call 811 or visit: <https://811.novascotia.ca>

Prepared by: Cardiac Services
Designed and Managed by: Nova Scotia Health Library Services

WG85-0698 © July 2025 Nova Scotia Health Authority
To be reviewed July 2028 or sooner, if needed.
Learn more: <https://library.nshealth.ca/patient-education-resources>