

Care After Sedation

Aussi disponible en français :

Précautions à prendre après une sédation (FF85-1679)



www.nshealth.ca

Care After Sedation

- Sedation is a treatment given to lower your pain during a procedure.
- You will be given medication through an intravenous (I.V.) injected into your vein with a needle. This is done by an Emergency Department (ED) staff member.
- The medication will make you more comfortable and relaxed. You may feel sleepy or drowsy during your procedure, and will feel less pain or no pain at all. Also, you may not remember parts of your procedure because of the medication.
- It is important for you and your caregivers at home to understand these instructions and to follow them when you leave the ED.

The medication you were given is:

- The effects of this medication should not last for more than 24 hours (1 day).

Instructions

- A responsible adult should drive you home and stay with you for 24 hours in case you need help. You should rest during these 24 hours.
- **Do not** walk or take the bus home. You may take a taxi home if you go with a responsible adult who can help you.
- While you are in the ED, you may get a prescription for medication to take at home. Talk about your prescription with your pharmacist.
- You may have nausea (feel sick to your stomach) after you have sedation. If so, have small amounts of clear liquids often, like:
 - › Flat (stirred) ginger ale or cola
 - › Broth
 - › Water
 - › Gatorade®
 - › Jell-O®
 - › Tea **without milk**
- Once your stomach feels better, slowly start having solid food again.
- **Do not drive a car, take part in dangerous activities, or sign any important documents.** Having sedation can impair (change) your judgement.

Medications

The checked boxes are your instructions.

☐ Medications advised for you:

☐ Continue your usual medications:

Follow-up instructions

- If you need follow-up care, ED staff will tell you where and when.
- You may need to go to a clinic at the hospital (like the cardiology clinic or the orthopedic clinic), or to your primary health care provider (family doctor or nurse practitioner). This will depend on the type of procedure you had.
- Follow-up care is usually within 7 days (1 week) of your procedure.

Go to the nearest Emergency Department right away if you have any of these symptoms:

- › Shortness of breath, chest pain, or palpitations (fluttering or racing feeling in your chest)
- › Bleeding that will not stop even after using a dressing
- › Pain and swelling that does not get better after pain medication and raising the injured part (if the pain is in your arm or leg)
- › Fever (temperature above 38 °C/100.4 °F)
- › Vomiting (throwing up) that does not stop or get better
- › You cannot keep liquids down
- › Feeling drowsy or unsteady on your feet 24 hours after you leave the ED

[illegible]

Looking for more health information?

Find this pamphlet and all our patient resources here: <https://library.nshealth.ca/PatientEducation>

Contact your local public library for books, videos, magazines, and other resources.

For more information, go to <http://library.novascotia.ca>

Connect with a registered nurse in Nova Scotia any time: call 811 or visit <https://811.novascotia.ca>

Learn about other programs and services in your community: call 211 or visit <http://ns.211.ca>

Nova Scotia Health promotes a smoke-free, vape-free, and scent-free environment.

Please do not use perfumed products. Thank you!

www.nshealth.ca

Prepared by: Emergency Services

Designed by: Nova Scotia Health Library Services

The information in this pamphlet is for informational and educational purposes only.

The information is not intended to be and does not constitute health care or medical advice.

If you have any questions, please ask your health care provider.

WA85-0121 © June 2023 Nova Scotia Health Authority

The information in this pamphlet is to be updated every 3 years or as needed.